

2022

Bat Assessment: Burton Park, Leopardstown Rd., Co. Dublin.



Dr Tina Aughney
Bat Eco Services

Bat Eco Services, Ulex House, Drumheel, Lisduff, Virginia, Co. Cavan. A82 XW62.

Licensed Bat Specialist: Dr Tina Aughney (tina@batecoservices.com, 086 4049468)

NPWS licence C13/2020 (Licence to handle bats, expires 31st December 2022)

NPWS licence 08/2020 (Licence to photograph/film bats, expires 31st December 2022)

NPWS licence DER/BAT 2022-36 (Survey licence, expires 24th March 2025).

Statement of Authority: Dr Aughney has worked as a Bat Specialist since 2000 and has undertaken extensive survey work for all Irish bat species including large scale development projects, road schemes, residential developments, wind farm developments and smaller projects in relation to building renovation or habitat enhancement. She is a monitoring co-ordinator and trainer for Bat Conservation Ireland. She is a co-author of the 2014 publication *Irish Bats in the 21st Century*. This book received the 2015 CIEEM award for Information Sharing. Dr Aughney is a contributing author for the Atlas of Mammals in Ireland 2010-2015.

All analysis and reporting is completed by Dr Tina Aughney. Data collected and surveying is completed with the assistance of a trained field assistant.

Mr. Shaun Boyle (Field Assistant) NPWS licence DER/BAT 2022-37 (Survey licence, expires 24th March 2025).

Client: Traynor Environmental on behalf of DLR Co. Co..

Project Name & Location: Burton Park, Leopardstown Road. Eircode D18 EK73.

Report Revision History

Date of Issue	Draft Number	Issued To (process of issuing)
19 th September 2022	Draft 1	By email to Traynor Environmental
26 th September 2022	Final	By email to Traynor Environmental
18 th October 2022	Separate Reports	By email to Traynor Environmental

Purpose

This document has been prepared as a Report for Traynor Environmental. Only the most up to-date report should be consulted. All previous drafts/reports are deemed redundant in relation to the named site.

Bat Eco Service accepts no responsibility or liability for any use that is made of this document other than by the client for the purposes for which it was originally commissioned and prepared.

Carbon Footprint Policy

It is the policy of Bat Eco Services to provide documentation digitally in order to reduce carbon footprint. Printing of reports etc. is avoided, where possible.

Bat Record Submission Policy

It is the policy of Bat Eco Services to submit all bat records to Bat Conservation Ireland database one year post-surveying. This is to ensure that a high level bat database is available for future desktop reviews. This action will be automatically undertaken unless otherwise requested, where there is genuine justification.

Executive Summary

Project Name & Location: Burton Park, Leopardstown Road. Eircode D18 EK73.

Proposed work: Upgrade of facilities.

Bat Survey Results - Summary

Bat Species	Roosts	Foraging	Commuting
Common pipistrelle <i>Pipistrellus pipistrellus</i>			√
Soprano pipistrelle <i>Pipistrellus pygmaeus</i>			√
Nathusius' pipistrelle <i>Pipistrellus nathusii</i>			
Leisler's bat <i>Nyctalus leisleri</i>			
Brown long-eared bat <i>Plecotus auritus</i>			
Daubenton's bat <i>Myotis daubentonii</i>			
Natterer's bat <i>Myotis nattereri</i>			
Whiskered bat <i>Myotis mystacinus</i>			
Lesser horseshoe bat <i>Rhinolophus hipposideros</i>			

Bat Survey Duties Completed (Indicated by red shading)

Tree PBR Survey	<input checked="" type="checkbox"/>	Daytime Building Inspection	<input checked="" type="checkbox"/>
Static Detector Survey	<input type="checkbox"/>	Daytime Bridge Inspection	<input type="checkbox"/>
Dusk Bat Survey	<input checked="" type="checkbox"/>	Dawn Bat Survey	<input type="checkbox"/>
Walking Transect	<input type="checkbox"/>	Driving Transect	<input type="checkbox"/>
Trapping / Mist Netting	<input type="checkbox"/>	IR Camcorder filming	<input type="checkbox"/>
Endoscope Inspection	<input type="checkbox"/>	Other	<input type="checkbox"/>

Citation: Bat Eco Services (2022) Bat Assessment: Burton Park, Leopardstown Road. Eircode D18 EK73. Unpublished report prepared for Traynor Environmental.

Contents

1. Introduction	4
1.1 Relevant Legislation & Bat Species Status in Ireland	4
1.1.1 Irish Statutory Provisions	4
1.1.2 EU Legislation	4
1.1.3 IUCN Red Lists	5
1.1.4 Irish Red List - Mammals	5
1.1.5 Irish Bat Species	5
1.2 Relevant Guidance Documents	7
1.2.1 Bat Survey Requirements & Timing	8
1.2.2 Evaluation & Assessment Criteria	11
1.2.3 Bat Mitigation Measures	16
1.3 Project Description	22
1.3.1 Site Locations	22
2. Bat Survey Methodology	23
2.1 Daytime Inspections	23
2.1.1 Building & Structure Inspection	23
2.1.2 Tree Potential Bat Roost (PBRs) Inspection	24
2.2 Night-time Bat Detector Surveys	25
2.2.1 Dusk Bat Surveys	25
2.3 Desktop Review	25
2.3.1 Bat Conservation Ireland Database	25
3. Bat Survey Results	27
3.1 Daytime Inspections	27
3.1.1 Building & Structure Inspection	27
3.1.2 Tree Potential Bat Roost (PBRs) Inspection	27
3.2 Night-time Bat Detector Surveys	27
3.2.1 Dusk Survey Site	27
3.3 Desktop Review	27
3.3.1 Bat Conservation Ireland Database	27
3.4 Survey Effort, Constraints & Survey Assessment	28
4. Bat Ecological Evaluation	29
4.1 Bat Species Recorded & Sensitivity	29
5. Impact Assessment & Mitigation	30
5.1 Assessment of Potential Impact - Foraging & Commuting Habitats	30
5.2 Assessment of Potential Impact - Construction & Operation of Development	30
5.3 Overall Assessment of Potential Impact	30
5.4 Bat Mitigation Measures	30
5.4.1 Lighting Plan	30
6. Survey Conclusions	31
7. Bibliography	32
8. Appendices	36
8.1 Appendix 1 – Bat Boxes	36
8.2 Appendix 2 Bat Assessment Tables	37

1. Introduction

Bat Eco Services was commissioned by Traynor Environmental to undertake a bat survey of Burton Park, Leopardstown Road. Eircode D18 EK73 and this entailed daytime inspection and dusk survey.

1.1 Relevant Legislation & Bat Species Status in Ireland

1.1.1 Irish Statutory Provisions

A small number of animals and plants are protected under Irish legislation (Nelson, *et al.*, 2019). The principal statutory provisions for the protection of animal and plant species are under the Wildlife Act 1976 (as amended) and the European Communities (Birds and Natural Habitats) Regulations 2011, as amended. The Flora (Protection) Order 2015 (S.I. no. 356 of 2015) lists the plant species protected by Section 21 of the Wildlife Acts. See www.npws.ie/legislation for further information.

The codes used for national legislation are as follows:

- WA = Wildlife Act, 1976, Wildlife (Amendment) Act, 2000 and other relevant amendments
- FPO = Flora (Protection) Order, 2015 (S.I. No. 356 of 2015)

1.1.2 EU Legislation

The Birds Directive (Directive 2009/147/EC) and Habitats Directive (Council Directive 92/43/EEC) are the legislative instruments which are transposed into Irish law, *inter alia*, by the European Communities (Birds and Natural Habitats) Regulations 2011 (S.I. No. 477 of 2011) ('the 2011' Regulations), as amended.

The codes used for the Habitats Directive (Council Directive 92/43/EEC) are:

- Annex II Animal and plant species listed in Annex II
- Annex IV Animal and plant species listed in Annex IV
- Annex V Animal and plant species listed in Annex V

The main aim of the Habitats Directive is the conservation of biodiversity by requiring Member States to take measures to maintain or restore natural habitats and wild species listed on the Annexes to the Directive at a favourable conservation status. These annexes list habitats (Annex I) and species (Annexes II, IV and V) which are considered threatened in the EU territory. The listed habitats and species represent a considerable proportion of biodiversity in Ireland and the Directive itself is one of the most important pieces of legislation governing the conservation of biodiversity in Europe.

Under Article 11 of the Directive, each member state is obliged to undertake surveillance of the conservation status of the natural habitats and species in the Annexes and under Article 17, to report to the European Commission every six years on their status and on the implementation of the measures taken under the Directive. In April 2019, Ireland submitted the third assessment of conservation status for 59 habitats and 60 species. There are three volumes with the third listing details of the species assessed.

Article 12 of the Habitats Directive requires Member States to take measures for the establishment of a strict protection regime for animal species listed in Annex IV(a) of the Habitats Directive within the whole territory of Member States. Article 16 provides for derogation from these provisions under defined conditions. These provisions are implemented under Regulations 51 and 54 of the 2011 Regulations.

1.1.3 IUCN Red Lists

The International Union for the Conservation of Nature (IUCN) coordinates the Red Listing process at the global level, defining the categories so that they are standardised across all taxa. Red Lists are also produced at regional, national and subnational levels using the same IUCN categories (IUCN 2012, 2019). Since 2009, Red Lists have been produced for the island of Ireland by the National Parks and Wildlife Service (NPWS) and the Northern Ireland Environment Agency (NIEA) using these IUCN categories. To date, 13 Red Lists have been completed. The Red Lists are an assessment of the risk of extinction of each species and not just an assessment of their rarity. Threatened species are those species categorised as Critically Endangered, Endangered or Vulnerable (IUCN, 2019) – also commonly referred to as ‘Red Listed’.

1.1.4 Irish Red List - Mammals

Red Lists in Ireland refer to the whole island, i.e. including Northern Ireland, and so follow the guidelines for regional assessments (IUCN, 2012, 2019). The abbreviations used are as follows:

- RE Regionally Extinct
- CR Critically Endangered
- EN Endangered
- VU Vulnerable
- NT Near Threatened
- DD Data Deficient
- LC Least Concern
- NA Not Assessed
- NE Not Evaluated

There are 27 terrestrial mammals species in Ireland, which includes the nine resident bat species listed. The terrestrial mammal, according to Marnell *et al.*, 2019, list for Ireland consists of all terrestrial species native to Ireland or naturalised in Ireland before 1500. The IUCN Red List categories and criteria are used to assess that status of wildlife. This was recently completed for the terrestrial mammals of Ireland. Apart from the two following two mammal species (grey wolf *Canis lupus* (regionally extinct) and black rat *Rattus rattus* (Vulnerable)), the remaining 25 species were assessed as least concern in the most recent IUCN Red List publication by NPWS (Marnell *et al.*, 2019).

1.1.5 Irish Bat Species

All Irish bat species are protected under the Wildlife Act (1976) and Wildlife Amendment Acts (2000 and 2010). Also, the EC Directive on The Conservation of Natural habitats and of Wild Fauna and Flora (Habitats Directive 1992), seeks to protect rare species, including bats, and their habitats and requires that appropriate monitoring of populations be undertaken. All Irish bats are listed in Annex IV of the Habitats Directive and the lesser horseshoe bat *Rhinolophus hipposideros* is further listed under Annex II. Across Europe, they are further protected under the Convention on the Conservation of European Wildlife and Natural Habitats (Bern Convention 1982), which, in relation to bats, exists to conserve all species and their habitats. The Convention on the Conservation of Migratory Species of Wild Animals (Bonn Convention 1979, enacted 1983) was instigated to protect migrant species across all European boundaries. The Irish government has ratified both these conventions.

Also, under existing legislation, the destruction, alteration or evacuation of a known bat roost is an offence. The most recent guidance document is “Guidance document on the strict protection of animal species of Community interest un the Habitats Directive (Brussels, 12.10.2021 C(2021) 7391 final”.

Regulation 51(2) of the 2011 Regulations provides –

“(2) Notwithstanding any consent, statutory or otherwise, given to a person by a public authority or held by a person, except in accordance with a licence granted by the Minister under Regulation 54, a person who in respect of the species referred to in Part 1 of the First Schedule—

(a) deliberately captures or kills any specimen of these species in the wild, (b) deliberately disturbs these species particularly during the period of breeding, rearing, hibernation and migration,

(c) deliberately takes or destroys eggs of those species from the wild,

(d) damages or destroys a breeding site or resting place of such an animal, or

(e) keeps, transports, sells, exchanges, offers for sale or offers for exchange any specimen of these species taken in the wild, other than those taken legally as referred to in Article 12(2) of the Habitats Directive,

shall be guilty of an offence.”

The grant of planning permission does not permit the commission of any of the above acts or render the requirement for a derogation licence unnecessary in respect of any of those acts.

Any works interfering with bats and especially their roosts, may only be carried out under a derogation licence granted by National Parks and Wildlife Service (NPWS) pursuant to Regulation 54 of the European Communities (Birds and Natural Habitats) Regulations 2011 (which transposed the EU Habitats Directive into Irish law).

There are eleven recorded bat species in Ireland, nine of which are considered resident on the island. Eight resident bat species and one of the vagrant bat species are vesper bats and all vespertilionid bats have a tragus (cartilaginous structure inside the pinna of the ear). Vesper bats are distributed throughout the island. Nathusius' pipistrelle *Pipistrellus nathusii* is a recent addition while the Brandt's bat has only been recorded once to-date (Only record confirmed by DNA testing, all other records has not been genetically confirmed). The ninth resident species is the lesser horseshoe bat *Rhinolophus hipposideros*, which belongs to the Rhinolophidea and has a complex nose leaf structure on the face, distinguishing it from the vesper bats. This species' current distribution is confined to the western seaboard counties of Mayo, Galway, Clare, Limerick, Kerry and Cork. The eleventh bat species, the greater horseshoe bat, was only recorded for the first time in February 2013 in County Wexford and is therefore considered to be a vagrant species. A total of 41 SACs have been designated for the Annex II species lesser horseshoe bat (1303), of which nine have also been selected for the Annex I habitat 'Caves not open to the public' (8310).

Irish bat species list is presented in Table 1 along with their current status.

Table 1: Status of the Irish bat fauna (Marnell *et al.*, 2019).

Species: Common Name	Irish Status	European Status	Global Status
Resident Bat Species ^			
Daubenton's bat <i>Myotis daubentonii</i>	Least Concern	Least Concern	Least Concern
Whiskered bat <i>Myotis mystacinus</i>	Least Concern	Least Concern	Least Concern
Natterer's bat <i>Myotis nattereri</i>	Least Concern	Least Concern	Least Concern
Leisler's bat <i>Nyctalus leisleri</i>	Least Concern	Least Concern	Least Concern
Nathusius' pipistrelle <i>Pipistrellus nathusii</i>	Least Concern	Least Concern	Least Concern
Common pipistrelle <i>Pipistrellus pipistrellus</i>	Least Concern	Least Concern	Least Concern
Soprano pipistrelle <i>Pipistrellus pygmaeus</i>	Least Concern	Least Concern	Least Concern
Brown long-eared bat <i>Plecotus auritus</i>	Least Concern	Least Concern	Least Concern
Lesser horseshoe bat <i>Rhinolophus hipposideros</i>	Least Concern	Least Concern	Least Concern
Possible Vagrants ^			
Brandt's bat <i>Myotis brandtii</i>	Data deficient	Least Concern	Least Concern
Greater horseshoe bat <i>Rhinolophus ferrumequinum</i>	Data deficient	Near threatened	Near threatened

^ Roche *et al.*, 2014

1.2 Relevant Guidance Documents

This report will draw on guidelines already available in Europe and will use the following documents:

- National Roads Authority (2006) Best Practice Guidelines for the Conservation of Bats in the Planning of National Road Schemes
- Collins, J. (Editor) (2016) Bat Surveys for Professional Ecologists: Good Practice Guidelines (3rd edition). Bat Conservation Trust, London
- McAney, K. (2006) A conservation plan for Irish vesper bats, Irish Wildlife Manual No. 20 National Parks and Wildlife Service, Department of Environment, Heritage and Local Government, Dublin, Ireland.
- Marnell, F., Kelleher, C. & Mullen, E. (2022) Bat mitigation guidelines for Ireland v2. Irish Wildlife Manuals, No. 134. National Parks and Wildlife Service, Department of Housing, Local Government and Heritage, Ireland (Version 1: Kelleher & Marnell, 2006).
- The status of EU protected habitats and species in Ireland: Conservation status in Ireland of habitats and species listed in the European Council Directive on the Conservation of Habitats, Flora and Fauna 92/43/EEC. National Parks and Wildlife Service, Department of Environment, Heritage and Local Government.
- Bat Conservation Trust (2018) Bats and artificial lighting in the UK: bats and the built environment series. Guidance Note 08/2019. BCT, London.
- Guidance document on the strict protection of animal species of Community interest un the Habitats Directive (Brussels, 12.10.2021 C(2021) 7391 final.
- EPA (2017) Guidelines on the information to be contained in Environmental Impact Assessment Reports.

Collins (2016) is the principal document used to provide guidance in relation to bat survey effort required but the level of surveying is assessed on a case-by-case basis taking into consideration the historical bat records for the survey area, presence of built, structures and trees potentially suitable for roosting bats and the presence of suitable bat habitats for foraging and commuting. Additional reference is made to this document in relation to determining the value of buildings, trees etc. as bat roosts. The tables referred to from this document are described in the following section and in the section on methodology.

Marnell *et al.* (2022) is referred to for guidance in relation to survey guidance (timing and survey design), derogation licences and mitigation measures.

1.2.1 Bat Survey Requirements & Timing

With reference to Collins (2016) and Marnell *et al.* (2022), the information presented in this section is used to determine the bat survey requirements for the proposed development site. Collins (2016) provides a trigger list in relation to determining if a bat survey is required and this is presented Appendix 3 (Figure B) for reference. In addition, Chapter 2 of Collins (2016) discusses that a bat survey is required when proposed activities are likely to impact on bats and their habitats. The level of surveying is to be determined by the ecologist and these are influenced by the following criteria:

- Likelihood of bats being present;
- Type of proposed activities;
- Scale of proposed activities;
- Size, nature and complexity of the site;
- Species concerned;
- No. of individuals.

Collins (2016) also provides the following table detailing when different survey components should be undertaken.

Table 2.2 Recommended UK survey times for survey types described in these guidelines.												
Survey type	Month											
	J	F	M	A	M	J	J	A	S	O	N	D
Preliminary ecological appraisal - fieldwork												
Preliminary roost assessment - structures ^a												
Emergence/re-entry survey for maternity or summer roosts ^b												
Emergence/re-entry ^c survey for transitional roosts ^b												
Emergence survey for mating roosts ^b												
Hibernation survey - structures ^a												
Preliminary ground level roost assessment - trees ^d												
Potential roost feature (PRF) inspection survey - trees												
Ground level bat activity survey - transects and automated/static												
Pre-, during and post-hibernation - automated/static bat activity survey												
Swarming survey												
Back-tracking survey												
Trapping survey ^e												
Radio tagging and tracking survey ^e												

= optimal period
 = sub-optimal period
 = weather or location dependent (i.e. may not be suitable due to spring and autumn conditions in any one year or in more northerly latitudes). Note that October surveys are not acceptable in Scotland.

Figure 1a: Table 2.2 reproduced from Collins (2016).

1.2.1.1 Buildings

In Marnell *et al.* (2022), Table 3 (The applicability of survey methods) provides information on the type of surveys that can be undertaken according to the different seasons.

Marnell *et al.* (2022) states that it is more suitable to survey buildings in the summer months. The following is a summary of the principal points:

1. The presence of a significant bat roost (invariably a maternity roost) can normally be determined on a single visit at any time of year, provided that the entire structure is accessible and that any signs of bats have not been removed by others. However, a visit during the summer or autumn has the advantage that bats may be seen or heard.
2. Roosts used by a small number of bats, as opposed to maternity sites, can be particularly difficult to detect and may require extensive searching backed up (in summer) by bat detector surveys or emergence counts.
3. If the entire building is not accessible or signs of bats may have been removed by others, or by the weather, bat detector or exit count methodologies may be required to back up a limited search.

Table 3. The applicability of survey methods.

Season	Roost type	Inspection	Bat detectors and emergence counts
Spring (Mar – May)	Building	Suitable (signs, perhaps bats)	Limited, weather dependent
	Trees	Difficult (best for signs before leaves appear)	Rarely useful
	Underground	Suitable (signs only)	Static detectors may be useful
Summer (June- August)	Building	Suitable (signs and bats)	Suitable
	Trees	Difficult	Limited; use sunrise survey
	Underground	Suitable (signs only)	Rarely useful
Autumn (September –November)	Building	Suitable (signs and bats)	Limited, weather dependent
	Trees	Difficult	Rather limited weather dependent; use sunrise survey?
	Underground	Suitable (signs, perhaps bats)	Static detectors may be useful
Winter (December- February)	Building	Suitable (signs, perhaps bats))	Rarely useful
	Trees	Difficult (best for signs after leaves have gone)	Rarely useful
	Underground	Suitable (signs and bats)	Static detectors may be useful

Figure 1b: Table 3 reproduced from Marnell *et al.* (2022).

The following table is used to determine the level and timing of surveys for buildings/structures with reference to the surrounding habitat. Buildings are assessed to determine their suitability as a bat roost and are described using the parameters Negligible, Low, Medium or High suitability in view of Table 2 from Marnell *et al.* (2022). The level of suitability informs the level of surveying and timing of surveys required based on Table 7.3 of Collins, 2016 (Note: These two tables are presented in Appendix 1 but a summary is provided in the table below).

Table 2a: Building Bat Roost Classification System & Survey Effort (Adapted from Collins, 2016 and Marnell *et al.*, 2022).

Suitability Category	Description (examples of criteria)	Survey Effort (Timings)
Negligible	Building have no potential as a roost site Urban setting, heavily disturbed, building material unsuitable, building in poor condition etc.	No surveys required.
Low	Building has a low potential as a roost site. No evidence of bat usage (e.g. droppings)	One dusk or dawn survey.
Medium	Building with some suitable voids / crevices for roosting bats. Some evidence of bat usage Suitable foraging and commuting habitat present.	At least one survey in May to August, minimum of two surveys (one dusk and one dawn).
High	Building with many features deemed suitable for roosting bats. Evidence of bat usage. Largely undisturbed setting, rural, suitable foraging and commuting habitat, suitable roof void and building material.	At least two surveys in May to August, with a minimum of three surveys (at least one dusk survey and one dawn survey).

1.2.1.2 Trees

Marnell *et al.* (2022) recommends the following in relation to detecting roosts in trees:

- “The best time to carry out surveys for suitable cavities is between November and April, when the trunk and branches are not obscured by leaves. If inspection suggests that the tree has suitable cavities or roost sites, a bat detector survey at dusk or dawn during the summer may help to produce evidence of bats, though the nomadic nature of most tree-dwelling species means that the success rate is very low.
- It can also be difficult to pinpoint exactly which tree a bat emerged from. A dawn survey is more likely to be productive than a dusk one as swarming bats returning to the roost are much more visible than those leaving the roost. Because tree-dwelling bats move roosts frequently, a single bat-detector survey is unlikely to provide adequate evidence of the absence of bats in trees that contain a variety of suitable roosting places.
- Several dawn or dusk surveys spread over a period of several weeks from June to August will greatly increase the probability of detecting significant maternity roosts and is recommended where development proposals will involve the loss of multiple trees”.

As a consequence, the BTHK (2018) Potential Roost Features (PRFs) list and the classification system adapted from Collins (2016) is recommended as part of the daytime inspection of trees to determine their PBR or Potential Bat Roost value. Details of the methodology followed is presented in Section 3.2.2.

1.2.1.3 Underground Structures

Marnell *et al.* (2022) recommends the following in relation to underground structures:

1. Underground structures are used mainly for hibernation, so surveys should generally be carried out during the winter.

1.2.2 Evaluation & Assessment Criteria

Based on the information collected during the desktop studies and bat surveys, an ecological value is assigned to each bat species recorded based on its conservation status at different geographical scales (Table 2b). For example, a site may be of national ecological value for a given species if it supports a significant proportion (e.g. 5%) of the total national population of that species.

Table 2b: The six-level ecological valuation scheme used in the CIEM Guidelines (2016) Ecological Value

Ecological Value	Geographical Scale of Importance
International	International or European scale
National	The Republic of Ireland or the island of Ireland scale (depending on the bat species)
Regional	Province scale: Leinster
County	County scale: County Dublin
Local	Proposed development and immediate surroundings
Negligible	None, the feature is common and widespread

If bat roosts are recorded, their roost status is determined using Figure 20 from Marnell *et al.* (2022). This figure is presented below (Figure 1c). This figure is also used to determine the conservation significance of the roost in order to prepare appropriate bat mitigation measures.

Impacts on bats can arise from activities that may result in:

- Physical disturbance of bat roosts e.g. destruction or renovation of buildings
- Noise disturbance e.g. increase human presence, use of machinery etc.
- Lighting disturbance
- Loss of roosts e.g. destruction or renovation of buildings
- Modifications of commuting or foraging habitats
- Severance or fragmentation of commuting routes
- Loss of foraging habitats.

It is recognised that any development will have an impact on the receiving environment, but the significance of the impact will depend on the value of the ecological features that would be affected. Such ecological features will be those that are considered to be important and potentially affected by the proposed development.

The guidelines consulted recommend that the potential impacts of a proposed development on bats are assessed as early as possible in the design stage to determine any areas of conflicts. In particular the Table 4 (presented as Figure 1d below) and Figure 20 (presented as Figure 1c) from Marnell *et al.* (2022) are referenced during this process.


Low	Roost status	Mitigation/compensation requirement (depending on impact)
Conservation significance 	Feeding perches of common/rarer species	Flexibility over provision of bat-boxes, access to new buildings etc. No conditions about timing or monitoring
	Individual bats of common species	
	Small numbers of common species. Not a maternity site	Provision of new roost facilities where possible. Need not be exactly like-for-like, but should be suitable, based on species' requirements. Minimal timing constraints or monitoring requirements
	Feeding perches of Annex II species	
	Small numbers of rarer species. Not a maternity site	
	Hibernation sites for small numbers of common/rarer species	Timing constraints. More or less like-for-like replacement. Bats not to be left without a roost and must be given time to find the replacement. Monitoring for 2 years preferred.
	Maternity sites of common species	
	Maternity sites of rarer species	Timing constraints. Like-for-like replacement as a minimum. No destruction of former roost until replacement completed and usage demonstrated. Monitoring for at least 2 years.
	Significant hibernation sites for rarer/rarest species or all species assemblages	
	Sites meeting SAC guidelines	
Maternity sites of rarest species		
High		Oppose interference with existing roosts or seek improved roost provision. Timing constraints. No destruction of former roost until replacement completed and significant usage demonstrated. Monitoring for as long as possible.

Figure 20 Guidelines for proportionate mitigation. The definition of common, rare and rarest species requires regional interpretation.

Figure 1c: Figure 20 (p 46) Reproduced from Marnell *et al.* (2022).

Table 4 The scale of main impacts at the site level on bat populations. [NB This is a general guide only and does not take into account species differences. Medium impacts, in particular, depend on the care with which any mitigation is designed and implemented and could range between high and low.]

Roost type	Development effect	Scale of impact		
		Low	Medium	High
Maternity	Destruction			✓
	Isolation caused by fragmentation			✓
	Partial destruction; modification		✓	
	Temporary disturbance outside breeding season	✓		
	Post-development interference			✓
Major hibernation	Destruction			✓
	Isolation caused by fragmentation			✓
	Partial destruction; modification		✓	
	Temporary disturbance outside hibernation season	✓		
	Post-development interference			✓
Minor hibernation	Destruction			✓
	Isolation caused by fragmentation			✓
	Partial destruction, modification		✓	
	Modified management		✓	
	Temporary disturbance outside hibernation season	✓		
	Post-development interference		✓	
	Temporary destruction, then reinstatement	✓		
Mating	Destruction		✓	
	Isolation caused by fragmentation		✓	
	Partial destruction	✓		
	Modified management	✓		
	Temporary disturbance	✓		
	Post-development interference	✓		
	Temporary destruction, then reinstatement	✓		
Night roost	Destruction	✓		
	Isolation caused by fragmentation	✓		
	Partial destruction	✓		
	Modified management	✓		
	Temporary disturbance	✓		
	Post-development interference	✓		
	Temporary destruction, then reinstatement	✓		

Figure 1d: Table 4 (p 44) Reproduced from Marnell *et al.* (2022).

Different parameters are considered for the overall assessment of the potential impact(s) of a proposed development on local bat populations.

The overall impacts of the proposed project on local bat populations is assessed using the following criteria:

- Impact Quality using the parameters Positive, Neutral or Negative Impact (based on EPA, 2017)

Table 2c: Criteria for assessing impact quality based on EPA, 2017,

Quality of Effect	Criteria
Positive	A change which improves the quality of the environment (for example, by increasing species diversity; or the improving reproductive capacity of an ecosystem, or by removing nuisances or improving amenities).
Neutral	No effects or effects that are imperceptible, within normal bounds of variation or within the margin of forecasting error.
Negative	A change which reduces the quality of the environment (for example, lessening species diversity or diminishing the reproductive capacity of an ecosystem; or damaging health or property or by causing nuisance).

- Impact Significance of potential impact parameters on specific bat species in relation to particular elements (e.g. roosting sites, foraging area and commuting routes) are assessed with reference to the following:
 - o Table 4 of Marnell *et al.* (2022) (Figure 1a);
 - o the known ecology and distribution of the bat species in Ireland;
 - o bat survey results including type of roosts (if any recorded), pattern of bat usage of the survey area, level of bat activity recorded etc.
 - o and bat specialist experience.
- Impact Significance of the proposed development on local bat populations maybe determine, where applicable, using the parameters listed in Table 2d (based on EPA, 2017).

Table 2d: Criteria for assessing significance of effects based on EPA, 2017,

Significance of Effects	Definition
Imperceptible	An effect capable of measurement but without significant consequences.
Not significant	An effect which causes noticeable changes in the character of the environment but without significant consequences.
Slight	An effect which causes noticeable changes in the character of the environment without affecting its sensitivities.
Moderate	An effect that alters the character of the environment in a manner that is consistent with existing and emerging baseline trends.
Significant	An effect which, by its character, magnitude, duration or intensity alters a sensitive aspect of the environment.
Very Significant	An effect which, by its character, magnitude, duration or intensity significantly alters most of a sensitive aspect of the environment.
Profound	An effect which obliterates sensitive characteristics

The following terms will be used, where possible and applicable, when quantifying the duration of the potential effects (selected from EPA, 2017):

- Temporary – effects lasting less than a year
- Short-term – effects lasting 1 to 7 years
- Medium term – effects lasting 7 to 15 years
- Long term – effects lasting 15 to 60 years
- Permanent – effects lasting over 60 years
- Reversible – effects that can be undone, for example through remediation or restoration

1.2.3 Bat Mitigation Measures

1.2.3.1 Bats & Lighting

All European bat species, including Irish bat species, are nocturnal. Light levels as low as typical full moon levels, i.e. around 0.1 LUX, can alter the flight activity of bats (Voigt *et al.* 2018). Any level of artificial light above that of moonlight can mask the natural rhythms of lunar sky brightness and, thus, can disrupt patterns of foraging and mating and might, for instance, interfere with entrainment of the circadian system.

Artificial light pollution is an increasing global problem (Rich and Longcore, 2006) and Artificial light at night (ALAN) is considered a major threat to biodiversity, especially to nocturnal species. As urbanisation expands into the landscape, the degree of street lighting also expands. Its ecological impacts can have a profound affect the behaviour of nocturnal animals including impacts on reproductive behaviours, orientation, predator-prey interaction and competition among others, depending on the taxon and ecosystem in question (Longcore and Rich 2004). It is considered by Hölker *et al.* (2010) to be a key biodiversity threat to biodiversity conservation. In relation to bats, the potential impacts of artificial night lighting can result in habitat fragmentation (Hanski, 1998), delay in roost emergence (Downs *et al.*, 2003) and a reduction in prey items.

In the context of behavioural ecology, lights can work to attract or repel certain animals. Many groups of insects, including moths, lacewings, beetles, bugs, caddisflies, crane flies, midges, hoverflies and wasps, can be attracted to artificial light (Eisenbeis and Hassel 2000; Frank 1988; Kolligs 2000). Attraction depends on the spectrum of light. In the context of street lights, white (mercury vapour) lamps emit a white light that includes ultraviolet. High pressure sodium lights (yellow) emit some ultraviolet, while low pressure sodium lamps (orange) emit no ultraviolet light (e.g. Rydell 2006). As a result of the attractiveness of lights to aerial invertebrates, swarms of insects often occur in and around street lights and, particular bat species such as aerial insect predators, can exploit the swarming insects to their advantage. Such attraction can also take prey items away from dark zones where light sensitive species are foraging, thus reducing their likelihood of feeding effectively.

Rydell (2006) divides bats into four categories in terms of their characteristic behaviours at street lamps. The four categories are based on bat size, wing morphology and echolocation call characteristics which were highlighted by Norberg and Rayner (1987) to determine flight speed, manoeuvrability, and prey detection capabilities of bats. Rydell (2006) stated that the large, fast flying bats, which are confined to open airspace, fly high over lit areas and are rarely observed near ground level. None of these, typically large free-tailed bats (e.g. large species of the family Molossididae), are found in Ireland. The second category are the medium-sized fast flying species, including the *Nyctalus* species, which patrol the street well above the lights and can be seen occasionally as they dive for prey into the light cone. This group includes the Leisler's bat, which is found in Ireland. Rydell's third category describes the small but fast flying bats that are manoeuvrable enough to

forage around light posts or under the lights, and includes the small *Pipistrellus* species of the old world, three of which are found in Ireland. The fourth category includes broad-winged slow flyers, most of which are seldom or never observed at lights. Slow flying bat species may be more vulnerable to predation by diurnal birds of prey and this may restrict their exploitation of insects around artificially illuminated areas (e.g. Speakman 1991). There are also the concerns that some bat species are more light sensitive and therefore actively avoid lit up areas. This is particularly relevant for lesser horseshoe bats. Therefore from this, we can categorise the suite of Irish bats species as follows (please note that the sensitivity category is the author’s description):

Table 3: Potential light sensitivity of the Irish bat fauna using categories described by Rydell, 2006.

Species: Common Name	Rydell Category	Sensitivity
Daubenton’s bat <i>Myotis daubentonii</i>	Category 4	Light sensitive
Whiskered bat <i>Myotis mystacinus</i>	Category 4	Light sensitive
Natterer’s bat <i>Myotis nattereri</i>	Category 4	Light sensitive
Leisler’s bat <i>Nyctalus leisleri</i>	Category 2	Light tolerant
Nathusius’ pipistrelle <i>Pipistrellus nathusii</i>	Category 3	Semi-tolerant
Common pipistrelle <i>Pipistrellus pipistrellus</i>	Category 3	Semi-tolerant
Soprano pipistrelle <i>Pipistrellus pygmaeus</i>	Category 3	Semi-tolerant
Brown long-eared bat <i>Plecotus auritus</i>	Category 4	Light sensitive
Lesser horseshoe bat <i>Rhinolophus hipposideros</i>	Category 4	Light sensitive

The ability of different bat species to exploit insects gathered around street lights varies greatly. Gleaning species such as *Myotis* bats rarely forage around street lights (Rydell and Racey, 1995). The ecological effects of illuminating aquatic habitats are also poorly known. Moore *et al.* (2006) found that light levels in an urban lake, subject simply to sky glow and not direct illumination from lights, reached the same order of magnitude as full moonlight.

All European bat species, including Irish bat species, are nocturnal. As a consequence, the scientific literature provides evidence that artificial lighting does impacts on bats. The degree of impact depends on the light sensitivity of the bat species and the type of luminaire. Lesser horseshoe bats are light sensitive and therefore adversely effected by the presence of lighting in all aspects of their life strategies (e.g. foraging, commuting, drinking and roosting).

The potential impacts of street lighting can be summarised as follows:

- Attracting Prey Items

Lights can work to attract or repel certain animals. Many groups of insects can be attracted to artificial light and this attraction depends on the spectrum of light. As a result of the attractiveness of lights to aerial invertebrates, swarms of insects often occur in and around street lights. Such attraction can also take prey items away from dark zones where light sensitive species, such as lesser horseshoe bats, are foraging, thus reducing their likelihood of feeding effectively.

- Reducing Foraging Habitat

The research documents that there is less bat species diversity foraging in habitats lit up by artificial lighting. Only bat species considered to be light tolerant are generally able to exploit habitats with lighting present, but overall, all bat species activity tends to be less in lit up habitats compared to non-lit up habitats.

- Fragmenting The Landscape

Scientific evidence shows that lighting is a barrier to the movement of light sensitive bat species, such as lesser horseshoe bats. Light sensitive bat species will actively seek dark corridors to commute along and therefore the presence of lighting in commuting habitats will restrict their movement of such species in the landscape.

- Reducing Drinking Sites

There is increasing evidence that drinking sites for bats is an essential component for local bat population survival and that the presence of artificial lighting at waterbodies prevents bats from availing of this resource.

Lighting, including street lights come in an array of different types but for street lights they typically include High Pressure Sodium, Low Pressure Sodium, Mercury Vapour and the more modern Light Emitting Diodes (LED). An array of field-based research has been undertaken to document the potential impact of lighting on bat flight activity. LED lighting is predicted to constitute 70% of the outdoor and residential lighting markets by 2020. While the use of LEDs promotes energy and cost savings relative to traditional lighting technologies, little is known about the effects these broad-spectrum “white” lights will have on wildlife, human health, animal welfare, and disease transmission. As a consequence, a large array of research has been undertaken recently on the potential impact of LED on bats.

Stone *et al.* (2012) undertook research in relation to “Cool” LED street lights on an array of local bat species in England. Overall the presence of LED street lights had a significant negative impact on lesser horseshoe bats and *Myotis* spp. for all light treatments investigated while there was no sign impact of light treatment type on *Pipistrellus pygmaeus* (soprano pipistrelle – a common Irish bat species) or *Nyctalus* (Leisler’s bats is part of this bat family and is a common Irish bat species)/*Eptesicus* species. This research paper also documented behavioural changes for the different bat species. Lesser horseshoe bats and *Myotis* spp. did not avoid lights by flying along the other side of the hedge but altered their commuting behaviour altogether. It was concluded that LEDs can fragment commuting routes causing bats to alter their behaviour with potentially negative conservation consequences. Lesser horseshoe bat activity was significantly lower during high intensity treatment than medium, but at all treatment levels (even as low as 3.6 LUX), activity was significantly lower than unlit control (LUX level measurements were taken at 1.7m at the hedge below the light).

Russo *et al.* (2017) investigated the impact of LED lighting on drinking areas for bats in Italy. Drinking sites are considered to be important components for the survival of local bat populations. Drinking sites were illuminated with a portable LED outdoor light emitting (48 high-power LEDs generated a light intensity of 6480 lm (4000–4500 K) at 25°C, two peaks of relative luminous flux at 450 and 590 nm). *Plecotus auritus* (brown long-eared bat – resident in Ireland), *Pipistrellus pygmaeus* (soprano pipistrelle – resident in Ireland) and *Rhinolophus hipposideros* (lesser horseshoe bat – resident in Ireland) did not drink when troughs were illuminated.

Rowse *et al.* (2018) researched the impacts of LED lights (portable lights, 97W 4250K LED on 10m high poles) in England on local bat populations. Treatments were either 100% light intensity; dimmed

(using pulse width modulation) at 50% or 25% light intensity; and unlit. Sites were in suburban areas along busy roads but with vegetation and tree lines adjacent. High light levels (50% & 100% light treatments) increased activity of opportunistic *Pipistrellus pipistrellus* (common pipistrelle – resident in Ireland) but reduced activity of *Myotis* species group. Conversely 25% and unlit sites had no difference from each other. The research paper conclude that dimming could be an effective strategy to mitigate ecological impacts of street lights.

Wakefield *et al.* (2017) stated that an important factor to be aware of in relation to LED is the direction of the light projected. Therefore it is recommended that highly focused/shielded LEDs designed to filter out short wavelengths of light may should be used as they attract relatively fewer insects. Less insects attracted to street lights means less insects leaving dark zones where light sensitive bat species primarily feed.

Martin *et al.* (2021) showed that LED street lights lead to a reduction in the total number of insects captured with light traps in a wide range of families. Coleoptera and Lepidoptera orders were the most sensitive groups to ecological light pollution in the study area. The paper suggested that LED was the least attractive light system for most of the affected groups both because of its very little emitted short-wavelength light and because of its lower light intensity. They also concluded that reduction in insect attraction to LED could be even larger with current LED technologies emitting warmer lights, since other research showed that LED emitting “warmer white” colour light (3000 K) involves significantly lower attraction for insects than “colder white” LED (6000 K).

Wilson *et al.* (2021) investigate the impact of LED on biting insects and concluded because LED is highly malleable with regard to spectral composition, they can be tailored to decrease or increase insect catches, depending on situation. Therefore this design control of LED could greatly assist in reducing impact of street lighting on local bat populations.

Stone *et al.* (2015) reviewed the impacts of ALAN on bat roosts and flight paths in order to provide recommendations in relation to street lighting. The principal recommendations were to avoid lighting places where bats are present and to ensure that there are interconnected light exclusion zones and variable light regimes with reduced intensity of light in specific areas (e.g. important foraging and commuting habitats) as responses to street lighting may vary between species. It recommends that there should be a 'light threshold'.

1.2.3.1.1 Lighting Guidelines – Effective Mitigation Measures

As a consequence of this extensive amount of research there are two principal guideline documents available for best practice for effective mitigation relating to outdoor lighting.

EUROBATS (2018) guidelines recommends the following:

- ALAN should be strictly avoided, and artificial lighting should be installed only where and when necessary coupled with the following:
 - o Dynamic lighting schemes, where possible.
 - o Use a minimal number of lighting points and luminaires on low positions in relation to the ground for minimising light trespass to adjacent bat habitats or into the sky.
 - o Use focused light, e.g. by using LED or shielded luminaires which limit the light flux only to the required areas and prevent light trespass into adjacent bat habitats.
 - o Create screens, either by erecting walls or by planting hedgerows or trees, to prevent light trespass, e.g. from illuminated roads, to surrounding bat habitats.
 - o Exits of bat roosts and a buffer zone around them should be protected from direct or indirect lighting to preserve the natural circadian rhythm of bats.

This BCT (2018) guidelines provides a list of recommendations in relation to luminaire design, which is based on the extensive research completed to-date on the potential impact of lighting on bats, and therefore provides best practice mitigation measures. These recommendations are the basis of mitigation measures pertaining to bats listed in this report and are summarised as follows:

- All luminaires used should lack UV/IR elements to reduce impact.
- A warm white spectrum (<2700 Kelvins should be used to reduce the blue light component of the LED spectrum).
- Luminaires should have a peak wavelengths higher than 550nm to avoid the component of light most disturbing to bats.
- Only luminaires with an upward light ratio of 0% and with good optical control should be used.
- Luminaires should be mounted on the horizontal, i.e. no upward tilt.
- Column heights should be carefully considered to minimise light spill. The shortest column height allowed should be used where possible.
- Bollard lighting should be considered for pedestrian, parks and greenway areas, if deemed necessary.

1.2.3.2 Bat Box Schemes

Bat Boxes are frequently used as part of bat mitigation to retain local bat populations within an area proposed to be development. The NPWS Survey and Mitigation Guidelines (Marnell *et al.* 2022) considers that where roosts of low conservation significance (Figure 20, Marnell *et al.* (2022)) are to be lost due to a development, bat boxes may provide an appropriate form of mitigation and the effectiveness depends on the type of bat box provided, which should be appropriate to the bat species (Figure 1f).

Species	Summer/ maternity	Summer/non breeding	Hibernation*	Notes
<i>Rhinolophus hipposideros</i>	N/A	N/A	N/A	Horseshoe bats cannot use bat boxes
<i>Myotis daubentonii</i>	H	H		
<i>Myotis mystacinus</i>	H	H		
<i>Myotis nattereri</i>	H	?		
<i>Pipistrellus nathusii</i>	H	H		
<i>Pipistrellus pipistrellus</i>	C	C/H	C	H are rarely used as maternity roosts.
<i>Pipistrellus pygmaeus</i>	C	C/H	C	
<i>Nyctalus leisleri</i>	H	H	H?	
<i>Plecotus auritus</i>	H	H		Maternity roosts
Key				
* Large well-insulated hibernation boxes may be more successful				
N/A -not applicable; bat boxes should not be considered as replacement roosts				
H - tree hollow-type box, providing a void in which bats can cluster				
C - tree crevice-type box, with 25-35mm crevices				
? - few data on which to base an assessment				

Figure 1e: Table 7 (p 58) Reproduced from Marnell *et al.* (2022).

1.2.3.3 Seasonality of Bat Mitigation Measures

The NPWS Survey and Mitigation Guidelines (Marnell *et al.* 2022) provides best practice guidance in relation to the timing of bat mitigation measures. It states that the most common and effective method of avoiding potential harm to a bat is to carry out the work at an appropriate time of the year. The following table provides a summary of timings.

Table 5 Optimum season for works in different types of roosts.

Bat usage of site	Optimum period for carrying out works (some variation between species)
Maternity	1 st October – 1 st May
Summer (not a proven maternity site)	1 st September – 1 st May
Hibernation	1 st May – 1 st October
Mating/swarming	1 st November – 1 st August

Figure 1f: Table 5 (p 50) Reproduced from Marnell *et al.* (2022).

Timing of bat mitigation measures is relevant to the proposed tree felling of Potential Bat Roosts (PBRs). Felling is recommended outside the principal maternity season and during mild weather conditions (to avoid cold weather that would encourage bats to hibernate). This coupled with dusk/dawn surveys and additional daytime inspections is best practice to ensure that tree felling is completed without causing harm to potentially roosting bats. The preferred tree felling months also avoids the bird nesting season.

1.3 Project Description

1.3.1 Site Locations

The proposed works related to under DLR Co. Co. management and are located in County Dublin.

Burton Park, Leopardstown Road. Eircode D18 EK73 (Red Line & Yellow Line areas).



Figure 2: Survey area - Burton Park, Leopardstown Road. Eircode D18 EK73.

2. Bat Survey Methodology

2.1 Daytime Inspections

One purpose of daytime inspections is to determine the potential of bat roosts within the survey area. Due to the transient nature of bats and their seasonal life cycle, there are a number of different type of bat roosts. Where possible, one of the objectives of the surveys is to be able to identify the types of roosts present, if any. However, the determination of the type of roost present depends on the timing of the survey and the number of bat surveys completed. Consequently, the definition of roost types, in this report, will be based on the following:

Table 4a: Bat Roost Types (adapted from Collins 2016).

Roost Type	Definition	Time of Survey
Day Roost	A place where individual bats or small groups of males, rest or shelter in the daytime but are rarely found by night in the summer.	Anytime of the year
Night Roost	A place where bats rest or shelter in the night but are rarely found in the day. May be used by a single bat on occasion or it could be used regularly by the whole colony.	Anytime of the year
Feeding Roost	A place where individual bats or a few bats rest or feed during the night but are rarely present by day.	Anytime of the year
Transitional Roost	A place used by a few individuals or occasionally small groups for generally short periods of time on waking from hibernation or in the period prior to hibernation.	Outside the main maternity and hibernation periods.
Swarming Site	Where large numbers of males and females gather. Appear to be important mating sites.	Late summer and autumn
Mating Site	Where mating takes place.	Late summer and autumn
Maternity Site	Where female bats give birth and raise their young to independence.	Summer months
Hibernation Site	Where bats are found, either individually or in groups in the winter months. They have a constant cool temperature and humidity.	Winter months in cold weather conditions
Satellite Roost	An alternative roost found in close proximity to the main nursery colony and is used by a few individuals throughout the breeding season.	Summer months

2.1.1 Building & Structure Inspection

Structures, buildings and other likely places that may provide a roosting space for bats are inspected during the daytime for evidence of bat usage. Evidence of bat usage is in the form of actual bats (visible or audible), bat droppings, urine staining, grease marks (oily secretions from glands present on stonework) and claw marks. In addition, the presence of bat fly pupae (bat parasite) also indicated that bat usage of a crevice, for example, has occurred in the past. Inspections are undertaken visually

with the aid of a strong torch beam (LED Lenser P14.2) and endoscope (General DC5660A Wet / Dry Scope).

Buildings were assessed to determine their suitability as a bat roost (26th August 2022) and described using the parameters Negligible, Low, Medium or High suitability in view of Marnell *et al.* (2022). The level of suitability informs the level of surveying required.

Table 4b: Building Bat Roost Classification System & Survey Effort (Adapted from Collins, 2016 and Marnell *et al.*, 2022).

Suitability Category	Description (examples of criteria)	Survey Effort (Timings)
Negligible	Building have no potential as a roost site Urban setting, heavily disturbed, building material unsuitable, building in poor condition etc.	No surveys required.
Low	Building has a low potential as a roost site. No evidence of bat usage (e.g. droppings)	One dusk or dawn survey.
Medium	Building with some suitable voids / crevices for roosting bats. Some evidence of bat usage Suitable foraging and commuting habitat present.	At least one survey in May to August, minimum of two surveys (one dusk and one dawn).
High	Building with many features deemed suitable for roosting bats. Evidence of bat usage. Largely undisturbed setting, rural, suitable foraging and commuting habitat, suitable roof void and building material.	At least two surveys in May to August, with a minimum of three surveys (at least one dusk survey and one dawn survey).

2.1.2 Tree Potential Bat Roost (PBRs) Inspection

Trees that may provide a roosting space for bats were classified using the Bat Tree Habitat Key (BTHK, 2018) and the classification system adapted from Collins (2016). The Potential Roost Features (PRFs) listed in this guide were used to determine the PBR value of trees.

Trees identified as PBRs were inspected during the daytime (26th August 2022), where possible, for evidence of bat usage. Evidence of bat usage is in the form of actual bats (visible or audible), bat droppings, urine staining, grease marks (oily secretions from glands present on stonework) and claw marks. In addition, the presence of bat fly pupae (bat parasite) also indicated that bat usage of a crevice, for example, has occurred in the past.

Daytime inspections were undertaken of all of the trees within the proposed development site. These inspections followed the Phase 1 guidance (Collins, 2016) in order to make a list of trees within the proposed development site that may be suitable as roosting sites for bats. Inspections were undertaken visually, from the ground, with the aid of a strong torch beam (LED Lenser P14.2) during the daytime searching for PRFs.

Table 4c: Tree Bat Roost Category Classification System (adapted from Collins, 2016).

Tree Category	Description
1 High	Trees with multiple, highly suitable features (Potential Roosting Features = PRFs) capable of supporting larger roosts
2 Moderate	Trees with definite bat potential but supporting features (PRFs) suitable for use by individual bats;
3 Low	Trees have no obvious potential although the tree is of a size and age that elevated surveys may result in cracks or crevices being found or the tree supports some features (PRFs) which may have limited potential to support bats;
4 Negligible	Trees have no potential.

2.2 Night-time Bat Detector Surveys

2.2.1 Dusk Bat Surveys

Dusk Emergence Survey was completed on the 28th August 2022 from 10 minutes before sunset to 110 minutes post sunset and the surveyors position themselves(s) within the proposed development sites to determine if bats were roosting within buildings and adjacent trees.

The following equipment was used:

Surveyor 2: Bat Logger M2 Full Spectrum Bat Detector and Pettersson D200 Heterodyne Bat Detector.

2.3 Desktop Review

2.3.1 Bat Conservation Ireland Database

Bat Conservation Ireland acts as the central depository for bat records for the Republic of Ireland. Its' bat database is comprised of >60,000 bat records. The database primarily contains bat records from the following datasets:

- Irish Bat Monitoring Programme

The Irish Bat Monitoring Programme is comprised of four surveys (Car-based Bat Monitoring Scheme (2003-), All Ireland Daubenton's Bat Waterways Survey (2006-), Brow Long-eared Bat Roost Monitoring Scheme (2007-) and Lesser Horseshoe Bat Monitoring Scheme (1980s-). Apart from the latter survey, all monitoring data is stored on the BCIreland database.

- BATLAS 2020 & 2010

BCIreland has undertaken two all-Ireland species distribution surveys (2008-2009 for BATLAS 2010 and 2016-2019 for BATLAS 2020) of four target bat species (Common and soprano pipistrelle, Leisler's bats and Daubenton's bat).

- Ad Hoc Bat Records

Ad hoc bat records from national bat groups, ecological consultants and BCIreland members are also stored on the BCIreland database.

- Roost Records

These records are only report at a 1km level to protect the location of private dwellings and to protect such important bat records.

A 1km radius search was requested for O1964426108.

3. Bat Survey Results

3.1 Daytime Inspections

3.1.1 Building & Structure Inspection

The array of buildings within all three sites were inspected on the 26th August 2022. All buildings were deemed to have a low potential for bat roosting. This took into consideration the type of structures present, the habitats location in the vicinity and the urban setting of the three sites.

3.1.2 Tree Potential Bat Roost (PBRs) Inspection

There are no tall vegetation within the site considered to have a Potential Bat Roost (PBR) value.

3.2 Night-time Bat Detector Surveys

3.2.1 Dusk Survey Site

Bat detector surveys completed on 28th August 2022 (Dusk Survey - Weather conditions: 15oC, patchy cloud cover, calm and dry).

No bats were recorded emerging from buildings or trees within the survey area. The first bat species recorded was a common pipistrelle at 20:59 hrs. A total of six passes were recorded between 20:29 and 21:38 hrs. A single soprano pipistrelle was briefly recorded at 21:38 hrs. No foraging was detected within the survey area. This was the extent of the bat activity in this area and it is a Low level of bat activity.

3.3 Desktop Review

3.3.1 Bat Conservation Ireland Database

A 1km radius search was requested for O1964426108:

No bat records.

3.4 Survey Effort, Constraints & Survey Assessment

The following table details any Survey Constraints encountered and a summary of Scientific Assessment completed.

Table 9: Survey Effort, Constraints & Survey Assessment Results.

Category	Discussion
Timing of surveys Surveying meets Collins, 2016 guidelines.	2022 Summer bat survey: 26 th and 28 th August 2022
Survey Type Full suite of surveys completed to ensure sufficient information was collated for bat assessment. Surveys completed according Collins, 2016 guidelines.	Bat Survey Duties Completed (Indicated by red shading) Tree PBR Survey <input checked="" type="checkbox"/> Daytime Building Inspection <input checked="" type="checkbox"/> Static Detector Survey <input type="checkbox"/> Daytime Bridge Inspection <input type="checkbox"/> Dusk Bat Survey <input checked="" type="checkbox"/> Dawn Bat Survey <input type="checkbox"/> Walking Transect <input type="checkbox"/> Driving Transect <input type="checkbox"/> Trapping/Mist Netting <input type="checkbox"/> IR Camcorder filming <input type="checkbox"/> Endoscope Inspection <input type="checkbox"/> Other (Thermal Imagery) <input type="checkbox"/>
Weather conditions	Suitable weather conditions for bat surveys
Survey Constraints	None
Survey effort	Summer bat survey: Daytime inspection – 1 hrs Dusk Survey – 2 hrs
Extent of survey area	Summer bat survey of proposed development sites.
Equipment	All in good working order.

The extent of the surveys undertaken has achieved to determine:

- Presence / absence of bat within the survey area;
- A bat species list for the survey area;
- Extent and pattern of usage by bats within the survey area.

It is therefore deemed that the Scientific Assessment completed is Appropriate in order to completed the aims of the bat survey.

4. Bat Ecological Evaluation

4.1 Bat Species Recorded & Sensitivity

Two species of bat was recorded within the survey areas: Soprano pipistrelle and common pipistrelle.

No bat roosts were recorded within the survey area.

A low level of bat activity was recorded for all bat species detected within the proposed survey areas.

Soprano pipistrelle

- Soprano pipistrelle is an Annex IV bat species under the EU Habitats Directive. The status of this bat species is listed as Least Concern. The national soprano pipistrelle population is considered to be significantly increasing trend (Aughney *et al.*, 2021).
- The modelled Core Area for soprano pipistrelle is a relatively large area that covers much of the island of Ireland (62,020km²). The Bat Conservation Ireland Irish Landscape Model indicated that the soprano pipistrelle selects areas with broadleaf woodland, riparian habitats and low density urbanisation (Roche *et al.*, 2014).

Common pipistrelle

- Common pipistrelle is an Annex IV bat species under the EU Habitats Directive. The status of this bat species is listed as Least Concern. The national common pipistrelle population is considered to be significantly increasing trend (Aughney *et al.*, 2021).
- The modelled Core Area for common pipistrelle is a relatively large area that covers much of the island of Ireland (56,485km²). The Bat Conservation Ireland Irish Landscape Model indicated that the Common pipistrelle selects areas with broadleaf woodland, riparian habitats and low density urbanization (<30%) (Roche *et al.*, 2014).

No Annex II bat species are known to occur in County Dublin (i.e. lesser horseshoe bat) and were not recorded within the survey.

5. Impact Assessment & Mitigation

Two bat species were recorded within the proposed development site and this represents 2 of the 8 resident bat species known for County Dublin. In addition, the level of bat activity within the proposed development site is considered to be Low for all of the bat species recorded during the bat surveys. Therefore it is deemed that the proposed development area is principally used as commuting routes for bats.

5.1 Assessment of Potential Impact - Foraging & Commuting Habitats

There is habitat within the proposed development sites suitable for commuting bats. The proposed development will not impact on commuting habitat for local bat populations.

5.2 Assessment of Potential Impact - Construction & Operation of Development

The construction of the proposed development works will potentially increase the degree of light. However, guidelines have been provided to reduce this potential impact.

5.3 Overall Assessment of Potential Impact -

The proposed development may result in the following:

- An increase in and light levels during construction.

Therefore the impact assessment is as follows:

- Habitat loss (potential foraging/ commuting habitat) effects on all bat species are assessed as **Imperceptible Negative Effects**.
- Disturbance and/or displacement effects on all bat species during the construction phase are assessed as **Short-term Imperceptible Negative Effect**

5.4 Bat Mitigation Measures

In order to reduce the potential negative impact of the proposed development on local bat populations, the following mitigation measures are recommended to be fully implemented. The Bat Mitigation Guidelines (Marnell *et al.* (2022) are the principal guidance in relation to bat mitigation in Ireland and therefore for this report.

Additional mitigation measures are also made in relation to lighting (BCT guidelines, 2018) and landscaping to further reduce the potential impact of the proposed development.

5.4.1 Lighting Plan

Bats are light sensitive bats species, hence their nocturnal activities. The three bat species recorded commuting and foraging within the survey area are Light Tolerant or Semi-tolerant bat species while the fourth bat species is Light Sensitive. Therefore, it is still important that strict lighting guidelines are required to reduce the potential impact of the proposed developments on local bat populations as standard best practice.

Luminaire design is extremely important to achieve an appropriate lighting regime. Luminaires come in a myriad of different styles, applications and specifications which a lighting professional can help to select. The following should be considered when choosing luminaires. This is taken from the most recent BCT Lighting Guidelines (BCT, 2018).

- All luminaires used will lack UV/IR elements to reduce impact.

- LED luminaires will be used due to the fact that they are highly directional, lower intensity, good colour rendition and dimming capability.
- A warm white spectrum (<2700 Kelvins will be used to reduce the blue light component of the LED spectrum).
- Luminaires will feature peak wavelengths higher than 550nm to avoid the component of light most disturbing to bats.
- Column heights should be carefully considered to minimise light spill. The shortest column height allowed should be used where possible. For this proposed development scheme bollard lighting will be used.
- Only luminaires with an upward light ratio of 0% and with good optical control will be used.
- Luminaires will be mounted on the horizontal, i.e. no upward tilt.
- Any external security lighting will be set on motion-sensors and short (1min) timers. For this proposed development scheme there is no security lighting.
- As a last resort, accessories such as baffles, hoods or louvres will be used to reduce light spill and direct it only to where it is needed.

6. Survey Conclusions

Two bat species were recorded within the proposed development sites and this represents 2 of the 8 resident bat species known for County Dublin. In addition, the level of bat activity within the proposed development sites is considered to be Low for all of the bat species recorded during the bat surveys. Therefore it is deemed that the proposed development areas are principally used as commuting routes for bats.

Bat mitigation measures have been provided to reduce potential negative impacts in relation to in relation to outdoor lighting.

7. Bibliography

- Abbott, I. M., Butler, F. And Harrison, S. (2012) When flyways meet highways – the relative permeability of different motorway crossing sites to functionality diverse bat species. *Landscape and Urban Planning* 106 (4): 293-302.
- Abbott, I. M., Berthinessen, A., Stone, E., Booman, M., Melber, M. and Altringham, J. (2015) Bats and Roads, Chapter 5, pp/ 290-299. In: *Handbook of Road Ecology*. Editors: R. Van der Ree., D. J. Smidt and C. Grilo. Wiley Blackwell.
- Altringham, J. D. (2013) *British Bats*. Collins New Naturalist Library, Volume 93. Haper Collins, London.
- Altringham, J. And Kerth, G. (2016) Bats and Roads, Chapter 3. In: *Bats in the Anthropocene: Conservation of Bats in a Changing World*. Editors: C. C. Voigt and T. Kingston. Springer Open.
- Aughney, T., Roche, N., & Langton, S (2018) The Irish Bat Monitoring Programme 2015-2017. *Irish Wildlife Manuals*, No. 103. National Parks and Wildlife Service, Department of Cultural heritage and the Gaeltacht, Ireland.
- Aughney, T., Roche, N. and Langton, S. (2022) Irish Bat Monitoring Programme 2018-2021. *Irish Wildlife Manuals*, No. 137. National Parks and Wildlife Service, Department of Housing, Local Government and Heritage, Ireland.
- Barratt, E. M., Deauville, R., Burland, T. M., Bruford, M. W., Jones, G., Racey, P. A., & Wayne, R. K. (1997). DNA answers the call of pipistrelle bat species. *Nature* 387: 138 - 139.
- Bat Conservation Ireland (2015) BATLAS 2020 Pilot Project 2015: Volunteer Survey Manual. Version 01. www.batconservationireland.org.
- Bat Conservation Trust (2018) Bats and artificial lighting in the UK: bats and the built environment series. Guidance Note 08/2019. BCT, London.
- Bharddwaj, M., Soaner, K., Straka, T., Lahoz-Monfort, J., Lumsden, L. F. and van der Ree, R. (2017) Differential use of highway underpasses by bats. *Biological Conservation* 212: 22-28.
- Billington, G. E. & Norman, G. M. (1997). A report on the survey and conservation of bat roosts in bridges in Cumbria, Kendal. *English Nature*.
- BTHK (2018) *Bat Roosts in Trees – A Guide to Identification and Assessment for Tree-Care and Ecology Professionals*. Exeter: Pelagic Publishing.
- CIEEM (2016) *Guidelines for Ecological impact Assessment in the UK and Ireland: Terrestrial, Freshwater and Coastal (2nd Edition)*. CIEEM, Winchester.
- Collins, J. (ed.) (2016) *Bat Surveys for Professional Ecologists: Good Practice Guidelines (3rd Edition)*. The Bat Conservation Trust, London.
- Collins, J.H., Ross, A.J., Ferguson, J.A., Williams, C.D. & Langton, S.D. (2022) The implementation and effectiveness of bat roost mitigation and compensation measures for *Pipistrellus* and *Myotis* spp. and brown long-eared bat (*Plecotus auritus*) included in building development projects completed between 2006 and 2014 in England and Wales. *Conservation Evidence*: 17, 19-26.
- Convention on the Conservation of European Wildlife and Natural Habitats (Bern Convention) 1982.
- Convention on the Conservation of Migratory Species of Wild Animals (Bonn Convention) 1979.
- Dietz, C., Helversen, O. and Dietmar, N. (2011) *Bats of Britain, Europe & Northweat Africa*. A&C Black, London.

Downs, N.C., Beaton, V., Guest, J., Polanski, J., Robinson, S.L. and Racey, P.A. (2003) The effects of illuminating the roost entrance on the emergence behaviour of *Pipistrellus pygmaeus*. *Biological Conservation* 111, p. 247-252.

EC Directive on The Conservation of Natural habitats and of Wild Fauna and Flora (Habitats Directive) 1992.

Eisenbeis G and Hassel F. (2000). Zur Anziehung nachtaktiver Insekten durch Straßenlaternen – eine Studie kommunaler Beleuchtungseinrichtungen in der Agrarlandschaft Reinheßens Attraction of nocturnal insects to street lights – a study of municipal lighting systems in a rural area of Rheinhessen (Germany)]. *Natur und Landschaft* 75: 145–56.

Frank K.D. (1988). Impact of outdoor lighting on moths: an assessment. *J Lepidop Soc* 42: 63–93.

Gunnell, K., Grant, G. and Williams, C (2012) Landscape and urban design for bats and biodiversity. The Bat Conservation Trust, London.

Hanski, I. (1998) Metapopulation Dynamics. *Nature*, 396, 41-49.

Holker, F., Wolter, C., Perkin, E.K. & Tockner, K. (2010). Light pollution as a biodiversity threat. *Trends Ecol. Evol.* 25, 681–682. <https://doi.org/10.1016/j.tree.2010.09.007>.

Hundt, L. (2012) Bat Surveys: Good Practice Guidelines (2nd Edition). The Bat Conservation Trust, London.

Kelleher, C. & Marnell, F. (2006) Bat Mitigation Guidelines for Ireland. Irish Wildlife Manuals, No. 25. National Parks and Wildlife Service, Department of Environment, Heritage and Local Government, Dublin, Ireland.

Kolligs D. 2000. Ökologische Auswirkungen künstlicher Lichtquellen auf nachtaktive Insekten, insbesondere Schmetterlinge (Lepidoptera) [Ecological effects of artificial light sources on nocturnally active insects, in particular on moths (Lepidoptera)]. *Faunistisch-Ökologische Mitteilungen Suppl* 28: 1–136.

Lintott P. & Mathews F. (2018) *Reviewing the evidence on mitigation strategies for bats in buildings: informing best-practice for policy makers and practitioners*. CIEEM Commissioned Report

Longcore T. and Rich C. (2004). Ecological light pollution. *Frontiers in Ecology and Environment*. 2: 191-198.

Lundy, M.G., Montgomery, I.W., Roche, N. & Aughney, T. (2011). *Landscape Conservation for Irish Bats & Species Specific Roosting Characteristics* (Unpublished). Bat Conservation Ireland, Cavan, Ireland.

Lysaght, L. and Marnell, F. (eds) (2016) Atlas of Mammals in Ireland 2010-2015, National Biodiversity Data Centre, Waterford.

Marnell, F., Kingston, N. & Looney, D. (2009) *Ireland Red List No. 3: Terrestrial Mammals*, National Parks and Wildlife Service, Department of the Environment, Heritage and Local Government, Dublin, Ireland.

Marnell, F., Looney, D. & Lawton, C. (2019) Ireland Red List No. 12: Terrestrial Mammals. National Parks and Wildlife Service, Department of the Culture, Heritage and the Gaeltacht, Dublin, Ireland.

Marnell, F., Kelleher, C. & Mullen, E. (2022) Bat mitigation guidelines for Ireland v2. Irish Wildlife Manuals, No. 134. National Parks and Wildlife Service, Department of Housing, Local Government and Heritage, Ireland.

Martín, B.; Pérez, H.; Ferrer, M. Light-Emitting Diodes (LED): A Promising Street Light System to Reduce the Attraction to Light of Insects. *Diversity* 2021, 13, 89. <https://doi.org/10.3390/d13020089>.

Mathews, F., Roche, N., Aughney, T., Jones, N.M. Day, J., Baker, J. and Langton, S. (2015) Barriers and benefits: implications of artificial night-lighting for the distribution of common bats in Britain and Ireland. *Philosophical Transactions of the Royal Society of London B* 370 (1667), doi: 10.1098/rstb.2014.0124.

- McAney, K. (2006) A conservation plan for Irish vesper bats, Irish Wildlife Manual No. 20 National Parks and Wildlife Service, Department of Environment, Heritage and Local Government, Dublin, Ireland. McAney, K. (2014). An overview of *Rhinolophus hipposideros* in Ireland (1994-2014). *Vespertilio* **17**, 115–125.
- McAney, K., O'Mahony, C., Kelleher, C., Taylor, A. & Biggane, S. (2013). *The Lesser Horseshoe Bat in Ireland: Surveys by The Vincent Wildlife Trust*. Belfast, Northern Ireland: Irish Naturalists' Journal.
- Mullen, E. (2007). Brandt's Bat *Myotis brandtii* in Co. Wicklow. Irish Naturalists' Journal 28: 343.
- Norberg U.M. and Rayner J.M.V. (1987). Ecological morphology and flight in bats (Mammalia; Chiroptera): wing adaptations, flight performance, foraging strategy and echolocation. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*. **316**: 335-427.
- NPWS (2018) Conservation objectives supporting document – lesser horseshoe bat (*Rhinolophus hipposideros*) Version 1. Conservation Objectives Supporting Document Series. National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht, Dublin, Ireland
- O'Sullivan, P. (1994). *Bats in Ireland*. Special supplement to the Irish Naturalists' Journal.
- Rich, C. & Longcore, T. (eds). 2006 Ecological consequences of artificial night lighting. Washington, DC: Island Press
- Richardson, P. (2000). *Distribution atlas of bats in Britain and Ireland 1980 - 1999*. The Bat Conservation Trust, London, UK.
- Roche, N., Aughney, T. & Langton, S. (2015). *Lesser Horseshoe Bat: population trends and status of its roosting resource* (No. 85). , Irish Wildlife Manuals. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht, Dublin, Ireland.
- Roche, N., Langton, S. & Aughney, T. (2012). *Lesser Horseshoe Bat: Population, Trends and Threats 1986 to 2012* (Unpublished). Bat Conservation Ireland, Cavan, Ireland.
- Roche, N., Aughney, T., Marnell, F. & Lundy, M. (2014). *Irish Bats in the 21st Century*. Bat Conservation Ireland, Cavan, Ireland.
- Rowse EG, Harris S, Jones G. 2018 Effects of dimming light-emitting diode street lights on light-opportunistic and light-averse bats in suburban habitats. *R.Soc. open sci.* **5**: 180205. <http://dx.doi.org/10.1098/rsos.180205>
- Russ, J. (2012) *British Bat Calls: A guide to species identification*. Pelagic Publishing, Exeter.
- Russo, D., Cistrone, L., Libralato, N., Korine, C., Jones, G. & Ancillotto, L. (2017). Adverse effects of artificial illumination on bat drinking activity. *Anim. Conserv.* **20**, 492–501. <https://doi.org/10.1111/acv.12340>.
- Rydell J. (1992). Exploitation of insects around streetlamps by bats in Sweden. *Functional Ecology* **6**: 744-750.
- Rydell J. (2006). Bats and their insect prey at streetlights. In C. Rich and T. Longcore (eds.) *Ecological Consequences of Artificial Night Lighting*. 43-60.
- Rydell J. and Racey P.A. (1995). Street lamps and the feeding ecology of insectivorous bats. In P.A. Racey and S.M. Swift (eds.) *Ecology, evolution and behaviour of bats. Symposia of the Zoological Society of London*. **67** pp 291-307. Clarendon Press, Oxford.
- Schofield, H. (2008). *The Lesser Horseshoe Bat Conservation Handbook*. Herefordshire, England: The Vincent Wildlife Trust.
- Speakman, J.R. (1991) Why do insectivorous bats in Britain not fly in daylight more frequently? *Funct. Ecol.* **5**, 518–524.

- Stebbing, R. E. & Walsh, S. T. (1991) *Bat Boxes: A guide to the history, function, construction and use in the conservation of bats*. The Bat Conservation Trust, 1991.
- Stone, E., Jones, G. and Harris, S. (2009). Street lighting disturbs commuting bats. *Current Biology*, **19**: 1123-1127.
- Stone, E. L., Jones, G., and Harris, S. (2012). Conserving energy at a cost to biodiversity? Impacts of LED lighting on bats. *Global Change Biology* **18**, 2458–2465. doi:10.1111/j.1365-2486.2012.02705.x
- Stone EL, Harris S, Jones G. 2015 Impacts of artificial lighting on bats: a review of challenges and solutions. *Mammal. Biol.* **80**, 213–219. (doi:10.1016/j.mambio.2015.02.004)
- Svensson A.M. and Rydell J. (1998). Mercury vapour lamps interfere with bat defence of tympanate moths (*Operophtera* spp.; Geometridae). *Animal Behaviour* **55**: 223-226.
- Voigt C.C., Azam, C., Dekker, J., Feguson, J., Fritze, M., Gazaryan, S., Holker, F., Jones, G., Leader, N., Limpens, H.J.G.A., Mathews, F., Rydell, J., Schofield, H., Spoelstra, K., Zagamajster, M. (2018) Guidelines for consideration of bats in lighting projects. EUORBATS Publication Series No. 8. UNEP/EUROBATS Secretariat, Bonn.
- Wakefield, A., Broyles, M., Stone, E.L., Jones, G. & Harris, S. (2016). Experimentally comparing the attractiveness of domestic lights to insects: Do LEDs attract fewer insects than conventional light types? *Ecol. Evol.* **6**, 8028–8036. <https://doi.org/10.1002/ece3.2527>.
- Whilde, A. (1993). *Threatened mammals, birds, amphibians and fish in Ireland. Irish Red Data Book 2: Vertebrates*. Belfast: HMSO.
- Wildlife Act 1976 and Wildlife [Amendment] Act 2000. Government of Ireland.
- Wilson, R., Wakefield, A., Roberts, N. and Jones, G. (2021) Artificial light and biting flies: the parallel development of attractive light traps and unattractive domestic lights. *Parasite & Vectors*. <https://doi.org/10.1186/s13071-020-04530-3>.
- Zeale, M.R.K., Stone, E.L., Zeale, E., Browne, W.J., Harris, S. & Jones, G. (2018). Experimentally manipulating light spectra reveals the importance of dark corridors for commuting bats. *Glob. Chang. Biol.* **24**, 5909–5918. <https://doi.org/10.1111/gcb.14462>.

8. Appendices

8.1 Appendix 1 – Bat Boxes

Examples of bat box design (self-cleaning boxes i.e. opened at the bottom to allow bat droppings to fall out).

- a) Woodcrete 1FF (Potential supplier - www.nhbs.com)



8.2 Appendix 2 Bat Assessment Tables

Table 4.1 Guidelines for assessing the potential suitability of proposed development sites for bats, based on the presence of habitat features within the landscape, to be applied using professional judgement.

Suitability	Description Roosting habitats	Commuting and foraging habitats
Negligible	Negligible habitat features on site likely to be used by roosting bats.	Negligible habitat features on site likely to be used by commuting or foraging bats.
Low	<p>A structure with one or more potential roost sites that could be used by individual bats opportunistically. However, these potential roost sites do not provide enough space, shelter, protection, appropriate conditions^a and/or suitable surrounding habitat to be used on a regular basis or by larger numbers of bats (i.e. unlikely to be suitable for maternity or hibernation^b).</p> <p>A tree of sufficient size and age to contain PRFs but with none seen from the ground or features seen with only very limited roosting potential.^c</p>	<p>Habitat that could be used by small numbers of commuting bats such as a gappy hedgerow or unvegetated stream, but isolated, i.e. not very well connected to the surrounding landscape by other habitat.</p> <p>Suitable, but isolated habitat that could be used by small numbers of foraging bats such as a lone tree (not in a parkland situation) or a patch of scrub.</p>
Moderate	A structure or tree with one or more potential roost sites that could be used by bats due to their size, shelter, protection, conditions ^a and surrounding habitat but unlikely to support a roost of high conservation status (with respect to roost type only – the assessments in this table are made irrespective of species conservation status, which is established after presence is confirmed).	<p>Continuous habitat connected to the wider landscape that could be used by bats for commuting such as lines of trees and scrub or linked back gardens.</p> <p>Habitat that is connected to the wider landscape that could be used by bats for foraging such as trees, scrub, grassland or water.</p>
High	A structure or tree with one or more potential roost sites that are obviously suitable for use by larger numbers of bats on a more regular basis and potentially for longer periods of time due to their size, shelter, protection, conditions ^a and surrounding habitat.	<p>Continuous, high-quality habitat that is well connected to the wider landscape that is likely to be used regularly by commuting bats such as river valleys, streams, hedgerows, lines of trees and woodland edge.</p> <p>High-quality habitat that is well connected to the wider landscape that is likely to be used regularly by foraging bats such as broadleaved woodland, tree-lined watercourses and grazed parkland.</p> <p>Site is close to and connected to known roosts.</p>

^a For example, in terms of temperature, humidity, height above ground level, light levels or levels of disturbance.

^b Evidence from the Netherlands shows mass swarming events of common pipistrelle bats in the autumn followed by mass hibernation in a diverse range of building types in urban environments (Korsten *et al.*, 2015). This phenomenon requires some research in the UK but ecologists should be aware of the potential for larger numbers of this species to be present during the autumn and winter in large buildings in highly urbanised environments.

^c This system of categorisation aligns with BS 8596:2015 Surveying for bats in trees and woodland (BSI, 2015).

Figure A: Table 4.1 (p 35) Reproduced from Collins (2016).

(1) Conversion, modification, demolition or removal of buildings (including hotels, schools, hospitals, churches, commercial premises and derelict buildings) which are:

- agricultural buildings (e.g. farmhouses, barns and outbuildings) of traditional brick or stone construction and/or with exposed wooden beams;
- buildings with weather boarding and/or hanging tiles that are within 200m of woodland and/or water;
- pre-1960 detached buildings and structures within 200m of woodland and/or water;
- pre-1914 buildings within 400m of woodland and/or water;
- pre-1914 buildings with gable ends or slate roofs, regardless of location;
- located within, or immediately adjacent to woodland and/or immediately adjacent to water;
- Dutch barns or livestock buildings with a single skin roof and board-and-gap or Yorkshire boarding if, following a preliminary roost assessment, the site appears to be particularly suited to bats.

(2) Development affecting built structures:

- tunnels, mines, kilns, ice-houses, adits, military fortifications, air-raid shelters, cellars and similar underground ducts and structures; unused industrial chimneys that are unlined and brick/stone construction;
- bridge structures, aqueducts and viaducts (especially over water and wet ground).

(3) Floodlighting of:

- churches and listed buildings, green space (e.g. sports pitches) within 50m of woodland, water, field hedgerows or lines of trees with connectivity to woodland or water;
- any building meeting the criteria listed in (1) above.

(4) Felling, removal or lopping of:

- woodland;
- field hedgerows and/or lines of trees with connectivity to woodland or water bodies;
- old and veteran trees that are more than 100 years old;
- mature trees with obvious holes, cracks or cavities, or that are covered with mature ivy (including large dead trees).

(5) Proposals affecting water bodies:

- in or within 200m of rivers, streams, canals, lakes, reed beds or other aquatic habitats.

(6) Proposals located in or immediately adjacent to:

- quarries or gravel pits;
- natural cliff faces and rock outcrops with crevices or caves and swallets.

(7) Proposals for wind farm developments of multiple wind turbines and single wind turbines (depending on the size and location) (NE TIN 051 – undergoing updates at the time of writing).

(8) All proposals in sites where bats are known to be present¹

This may include proposed development affecting any type of buildings, structures, feature or location.

Notes:

1. Where sites are of international importance to bats, they may be designated as SACs. Developers of large sites 5–10km away from such SACs may be required to undertake a HRA.

Figure B: Reproduced from Collins (2016) – page 13.

Table 2 Factors affecting the probability of bats being present.

Factors affecting the probability of a building being used by bats in summer	
Increased probability	Disused or little used; largely undisturbed Large roof void with unobstructed flying spaces Large dimension roof timbers with cracks, joints and holes Uneven roof covering with gaps, though not too draughty Entrances that bats can fly in through Hanging tiles or wood cladding, especially on south-facing walls Rural setting Close to woodland and/or water Pre-20 th century or early 20 th century construction Roof warmed by the sun Within the distribution area of horseshoe bats
Decreased probability	Highly urbanised area with few feeding places Small or cluttered roof void (esp. for brown long-eared bat) Heavily disturbed Modern construction with few gaps around soffits or eaves (but be aware these may be used by pipistrelles in particular) Prefabricated with steel and sheet materials Active industrial premises Roof shaded from the sun
Factors affecting the probability of trees being used by roosting bats	
Increased probability	In ancient woodland or parkland Large trees with complex growth form Species that typically form cavities, such as beech, willow, oak or ash Visible damage caused by rot, wind, lightning strike <i>etc.</i> Loose bark providing cavities
Decreased probability	Coniferous plantation with no specimen trees Young trees with simple growth form and little damage
Factors affecting the probability of underground sites being used by roosting bats	
Increased probability	Large enough to develop stable temperature in winter High humidity Undisturbed Close to woodland or water (but note that bats will also use upland sites) Many cracks and crevices suitable for bats
Decreased probability	Small and draughty Heavily disturbed In urbanised areas Smooth surfaces with few roosting opportunities

Figure C: Table 2 Reproduced from Marnell *et al.* (2022).